A PHEROMONE PRODUCT FOR CONTROL OF LEPIDOPTERAN PESTS OF STORED PRODUCTS BY MATING DISRUPTION

Blue dispenser without antioxidant

## **ACTIVE INGREDIENT**

Z-9, E-12-Tetradecadien-1-yl acetate	- 93 00%
OTHER INGREDIENTS	- 7.00%
TOTAL	- 100 00%

Each dispenser contains 750 mg of pheromone

## **KEEP OUT OF REACH OF CHILDREN**

## CAUTION

SEE BACK PANEL FOR FIRST AID STATEMENTS
READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

ChemTica USA 2912 Enterprise Blvd Durant, Oklahoma 74701

Not EPA registered: For trial use. Each dispenser contains a magnet for hanging lure.

## **CONTENTS:**

5 dispensers, each containing 750 mg Z-9, E-12-tetradecadien-1-yl acetate per dispenser

NET WEIGHT 1.78 ounces/50.39 grams

## CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Avoid breathing vapor. Remove contaminated clothes and wash before reuse. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

#### **FIRST AID**

## IF ON SKIN OR CLOTHING

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call poison control center or doctor for treatment advice.

## IF INHALED

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.

Call poison control center or doctor for treatment advice.

## IF IN EYES

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 (between 9.30 am and 7.30 pm eastern time) for emergency treatment advice.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## 1. SITES

This product is to be used for the control of lepidopteran pests of stored products in Apartments, Bakeries, Commercial & Industrial Structures, Day Care Facilities, Food/Feed Manufacturing Facilities, Homes, Hospitals & Nursing Homes, Hotels/Motels, Kennels, Livestock operations, Meat Packing & Food Processing Plants, Mills, Pet Stores, Restaurants, Retail Establishments, Schools, Supermarkets, Transportation Equipment (buses, boats, planes, trains, trucks) and Warehouses

## 2. PESTS:

Place dispensers every 20-40 feet at chest height along the perimeter of the area to be treated. Place additional dispensers on the interior support structures (target 1 dispenser every 500 - 1,500 ft <sup>2</sup>). Add new dispensers every 6 months [180 days] or when monitoring traps show increased captures. Dispensers may be left in place for 9-12 months to ensure all pheromone is utilized.

As promulgated at 40 CFR 180.1153 DO NOT EXCEED the rate of 3.5 grams pheromone per 1,000 square feet/year.

## 4. TIMING:

Deploy after detection of stored products moths in pheromone monitoring traps. Dispenser will emit effective amounts of pheromone for 6 months [180 days].

- NOTE:(i) This product impedes the mating of many species of stored product moths. If pest populations densities are excessively high, disruption of mating will be less effective. At high population densities insecticide applications are recommended to reduce populations to manageable levels.
- (iii) Monitoring of male stored products moths using pheromone traps is recommended in treated areas. As a general rule, pheromone trap monitoring of other pest species should be done so that timely intervention is possible when warranted.

## STORAGE AND DISPOSAL STATEMENTS

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original package in dark. Below 25°C/77°F. Refrigerate after opening.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Dispose of empty bag in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **CONDITIONS OF SALE AND WARRANTY**

A PHEROMONE PRODUCT FOR CONTROL OF LEPIDOPTERAN PESTS OF STORED PRODUCTS BY MATING DISRUPTION

Brown dispenser without antioxidant

## **ACTIVE INGREDIENT**

Z-9, E-12-Tetradecadien-1-yl acetate	- 93 00%
OTHER INGREDIENTS	. 7.00%
TOTAL	100.00%

Each dispenser contains 750 mg of pheromone

## **KEEP OUT OF REACH OF CHILDREN**

## CAUTION

SEE BACK PANEL FOR FIRST AID STATEMENTS READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

ChemTica USA 2912 Enterprise Blvd Durant, Oklahoma 74701

Not EPA registered: For trial use. Each dispenser contains a magnet for hanging lure.

## **CONTENTS:**

5 dispensers, each containing 750 mg Z-9, E-12-tetradecadien-1-yl acetate per dispenser

NET WEIGHT 1.80 ounces/51.17 grams

## CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Avoid breathing vapor. Remove contaminated clothes and wash before reuse. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

#### **FIRST AID**

## IF ON SKIN OR CLOTHING

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call poison control center or doctor for treatment advice.

## IF INHALED

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.

Call poison control center or doctor for treatment advice.

## IF IN EYES

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 (between 9.30 am and 7.30 pm eastern time) for emergency treatment advice.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## 1. SITES

This product is to be used for the control of lepidopteran pests of stored products in Apartments, Bakeries, Commercial & Industrial Structures, Day Care Facilities, Food/Feed Manufacturing Facilities, Homes, Hospitals & Nursing Homes, Hotels/Motels, Kennels, Livestock operations, Meat Packing & Food Processing Plants, Mills, Pet Stores, Restaurants, Retail Establishments, Schools, Supermarkets, Transportation Equipment (buses, boats, planes, trains, trucks) and Warehouses

## 2. PESTS:

Place dispensers every 20-40 feet at chest height along the perimeter of the area to be treated. Place additional dispensers on the interior support structures (target 1 dispenser every 500 - 1,500 ft <sup>2</sup>). Add new dispensers every 6 months [180 days] or when monitoring traps show increased captures. Dispensers may be left in place for 9-12 months to ensure all pheromone is utilized.

As promulgated at 40 CFR 180.1153 DO NOT EXCEED the rate of 3.5 grams pheromone per 1,000 square feet/year.

## 4. TIMING:

Deploy after detection of stored products moths in pheromone monitoring traps. Dispenser will emit effective amounts of pheromone for 6 months [180 days].

- NOTE:(i) This product impedes the mating of many species of stored product moths. If pest populations densities are excessively high, disruption of mating will be less effective. At high population densities insecticide applications are recommended to reduce populations to manageable levels.
- (iii) Monitoring of male stored products moths using pheromone traps is recommended in treated areas. As a general rule, pheromone trap monitoring of other pest species should be done so that timely intervention is possible when warranted.

## STORAGE AND DISPOSAL STATEMENTS

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original package in dark. Below 25°C/77°F. Refrigerate after opening.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Dispose of empty bag in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **CONDITIONS OF SALE AND WARRANTY**

A PHEROMONE PRODUCT FOR CONTROL OF LEPIDOPTERAN PESTS OF STORED PRODUCTS BY MATING DISRUPTION

Blue dispenser with antioxidant

## **ACTIVE INGREDIENT**

Z-9, E-12-Tetradecadien-1-yl acetate	87.00%
OTHER INGREDIENTS	
α-Tocopherol (Vitamin E)	6.00%
TOTAL	100.00%

Each dispenser contains 700 mg of pheromone

## **KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

SEE BACK PANEL FOR FIRST AID STATEMENTS
READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

ChemTica USA 2912 Enterprise Blvd Durant, Oklahoma 74701

Not EPA registered: For trial use. Each dispenser contains a magnet for hanging lure.

## **CONTENTS:**

5 dispensers, each containing 700 mg Z-9, E-12-tetradecadien-1-yl acetate per dispenser

NET WEIGHT 1.78 ounces/50.36 grams

## CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Avoid breathing vapor. Remove contaminated clothes and wash before reuse. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

#### **FIRST AID**

#### IF ON SKIN OR CLOTHING

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call poison control center or doctor for treatment advice.

## **IF INHALED**

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.

Call poison control center or doctor for treatment advice.

#### IF IN EYES

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 (between 9.30 am and 7.30 pm eastern time) for emergency treatment advice.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## 1. SITES

This product is to be used for the control of lepidopteran pests of stored products in Apartments, Bakeries, Commercial & Industrial Structures, Day Care Facilities, Food/Feed Manufacturing Facilities, Homes, Hospitals & Nursing Homes, Hotels/Motels, Kennels, Livestock operations, Meat Packing & Food Processing Plants, Mills, Pet Stores, Restaurants, Retail Establishments, Schools, Supermarkets, Transportation Equipment (buses, boats, planes, trains, trucks) and Warehouses

## 2. PESTS:

Place dispensers every 20-40 feet at chest height along the perimeter of the area to be treated. Place additional dispensers on the interior support structures (target 1 dispenser every 500 - 1,500 ft ²). Add new dispensers every 6 months [180 days] or when monitoring traps show increased captures. Dispensers may be left in place for 9-12 months to ensure all pheromone is utilized.

As promulgated at 40 CFR 180.1153 DO NOT EXCEED the rate of 3.5 grams pheromone per 1,000 square feet/year.

## 4. TIMING:

Deploy after detection of stored products moths in pheromone monitoring traps. Dispenser will emit effective amounts of pheromone for 6 months [180 days].

- NOTE:(i) This product impedes the mating of many species of stored product moths. If pest populations densities are excessively high, disruption of mating will be less effective. At high population densities insecticide applications are recommended to reduce populations to manageable levels.
- (iii) Monitoring of male stored products moths using pheromone traps is recommended in treated areas. As a general rule, pheromone trap monitoring of other pest species should be done so that timely intervention is possible when warranted.

## STORAGE AND DISPOSAL STATEMENTS

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original package in dark. Below 25°C/77°F. Refrigerate after opening.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Dispose of empty bag in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **CONDITIONS OF SALE AND WARRANTY**

A PHEROMONE PRODUCT FOR CONTROL OF LEPIDOPTERAN PESTS OF STORED PRODUCTS BY MATING DISRUPTION

Brown dispenser with antioxidant

## **ACTIVE INGREDIENT**

Z-9, E-12-Tetradecadien-1-yl acetate	87.00%
OTHER INGREDIENTS	
α-Tocopherol (Vitamin E)	6.00%
TOTAL	100.00%

Each dispenser contains 700 mg of pheromone

## **KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

SEE BACK PANEL FOR FIRST AID STATEMENTS
READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

ChemTica USA 2912 Enterprise Blvd Durant, Oklahoma 74701

Not EPA registered: For trial use. Each dispenser contains a magnet for hanging lure.

## **CONTENTS:**

5 dispensers, each containing 700 mg Z-9, E-12-tetradecadien-1-yl acetate per dispenser

NET WEIGHT 1.80 ounces/51.14 grams

## CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Avoid breathing vapor. Remove contaminated clothes and wash before reuse. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

#### **FIRST AID**

#### IF ON SKIN OR CLOTHING

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call poison control center or doctor for treatment advice.

## **IF INHALED**

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible.

Call poison control center or doctor for treatment advice.

#### IF IN EYES

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present after 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 (between 9.30 am and 7.30 pm eastern time) for emergency treatment advice.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## 1. SITES

This product is to be used for the control of lepidopteran pests of stored products in Apartments, Bakeries, Commercial & Industrial Structures, Day Care Facilities, Food/Feed Manufacturing Facilities, Homes, Hospitals & Nursing Homes, Hotels/Motels, Kennels, Livestock operations, Meat Packing & Food Processing Plants, Mills, Pet Stores, Restaurants, Retail Establishments, Schools, Supermarkets, Transportation Equipment (buses, boats, planes, trains, trucks) and Warehouses

## 2. PESTS:

Place dispensers every 20-40 feet at chest height along the perimeter of the area to be treated. Place additional dispensers on the interior support structures (target 1 dispenser every 500 - 1,500 ft ²). Add new dispensers every 6 months [180 days] or when monitoring traps show increased captures. Dispensers may be left in place for 9-12 months to ensure all pheromone is utilized.

As promulgated at 40 CFR 180.1153 DO NOT EXCEED the rate of 3.5 grams pheromone per 1,000 square feet/year.

## 4. TIMING:

Deploy after detection of stored products moths in pheromone monitoring traps. Dispenser will emit effective amounts of pheromone for 6 months [180 days].

- NOTE:(i) This product impedes the mating of many species of stored product moths. If pest populations densities are excessively high, disruption of mating will be less effective. At high population densities insecticide applications are recommended to reduce populations to manageable levels.
- (iii) Monitoring of male stored products moths using pheromone traps is recommended in treated areas. As a general rule, pheromone trap monitoring of other pest species should be done so that timely intervention is possible when warranted.

## STORAGE AND DISPOSAL STATEMENTS

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original package in dark. Below 25°C/77°F. Refrigerate after opening.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Dispose of empty bag in a sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **CONDITIONS OF SALE AND WARRANTY**